# Three-dimensional digital planning of class III decompensation with clear aligners: Hard and soft tissue augmentation with concomitant corticotomy to stretch the limits of safe orthodontic treatment

### Q6

*i*) The corrections made in this section will be reviewed and approved by a journal production editor.

Federico Brugnami<sup>a,\*</sup> fbrugnami@gmail.com, Simonetta Meuli<sup>b</sup> simonetta.meuli@fastwebnet.it, Alfonso Caiazzo<sup>4</sup> Alfonsocaiazzo7@gmail.com, Sergio Marrocco<sup>4</sup> marrocco1sergio@gmail.com, Domenico Scopelliti, <u>Head of</u>

Q1 <u>Department</u><sup>d</sup> scopelliti61@gmail.com

<sup>a</sup>Piazza dei Prati degli Strozi 21, 00195, Rome, Italy

<sup>b</sup>Visiting Professor Tor Vergata University of Rome Italy, Via Ridolfino Venuti 36, 00189, Rome, Italy

<sup>e</sup>Clinical Assistant Professor Department of Oral and Maxillofacial Surgery, MGSDM Boston University, Salerno Italy, Via Bottiglieri 32, 00035, Salerno, Italy

<sup>d</sup>Department of Oral Maxillofacial Surgery San Filippo Neri Hospital Rome Italy, Smile House Italy, Operation Smile Foundation, Via Trionfale 1023, 00100, Rome, Italy

Corresponding author. DDS Piazza dei Prati degli Strozzi 21, 00195, Rome, Italy.

### Abstract

**Purpose:** Three-dimensional diagnosis has shown that orthodontic therapy could potentially move the roots of the teeth outside the original bone structure. The purpose of these case studies was to test the possibility of obtaining correct three-dimensional tooth positioning with clear aligners, thereby modifying the periodontal structure accordingly, at the same time.

*Methods*: Regenerative Corticotomy (RC) was performed with clear aligners on ten adult patients (40 anterior teeth) with skeletal Class III malocclusion, for dental decompensation, prior to the orthognathic surgery. The CBCT examinations were performed before treatment (T0) and 1 year after orthognathic surgery (T1). The vertical and the horizontal hard tissue changes, the width of keratinized gingiva, the incisors proclination (IMPA) and the percentage of inclination compared to the planning were analyzed.

**Results :** The distance between the Cemento-Enamel Junction (CEJ) and the Bone Marginal Level (BML) decreased in average from  $5.5 \pm 3.2$  mm to  $1.39 \pm 0.53$  mm. The horizontal changes were at the 3 mm level  $1.42 \pm 0.5$  mm, at the 5 mm level  $1.98 \pm 0.66$  mm and at the 7 mm level  $2.70 \pm 0.87$  mm. The width of Keratinized gingiva changes were on average  $1.42 \pm 0.36$  at T0 and  $4.16 \pm 2.25$  at T1. All the changes were statistically significant with p < 0,05. The mean proclination based on IMPA values was +9.16 +-1.19°. The mean difference of the incisor's proclination compared to the digitally planned was  $-1 + 0.6^{\circ}$  (89.87 +- 6.46%).

**Conclusion :** Clear Aligners with three-dimensional digital planning seems to be reliable in controlling teeth movements in the pre-orthognathic decompensation phase. Regenerative Corticotomy seems to have the ability to improve the periodontal tissues despite proclination.

**Keywords**: Orthognatic surgery; Clear aligners; Regenerative corticotomy; PAOO; Digital planning; Orthodontics-periodontics

# **1** Introduction

Three-dimensional diagnosis demonstrates that orthodontic therapy can potentially move the roots of the teeth outside the original bony structure.<sup>1</sup> This may lead to several periodontal problems. Three-dimensional digital planning may help to foresee such possible movements. The cases whose objective of the orthodontic treatment is the skeletal Class III orthognathic decompensation, may displace the roots outside the original bony structure. This means that periodontal side effects, like resorption of the buccal plate and/or the thinning of the overlying gingiva, are highly probable.<sup>2</sup> Furthermore, a higher risk of gingival recession may occur and the presence of a rapid progression of periodontal disease in susceptible patients several years after completion of the treatment could be an eventuality.<sup>3</sup> Studies have shown that Regenerative Corticotomy (RC) (also known as: PAOO<sup>®</sup>, Periodontally Accelerated Osteogenic Orthodontics) can counteract the marginal bone resorption due to root movement outside the bony structure, both in conventional treatment<sup>4,5</sup> and in orthognathic cases.<sup>6</sup>

Authors have not focused on Clear Aligners (CA) nor on the associated effect of RC on soft tissue reactions. Ahn and colleagues, in 2016,<sup>7</sup> evaluated two groups of 15 patients undergoing Class III orthognathic correction. The decompensation was associated with Augmentation Corticotomy only in the Test group, resulting in a preservation or augmentation of the periodontal structures after the decompensation. This complies with the application of Augmented Corticotomy in orthodontic treatment in non-surgical cases.<sup>8</sup> Many studies have involved patients treated with brackets and wires. CA is a viable alternative to brackets and wires to correct many different malocclusions. Various authors have questioned the ability of CA in controlling tooth movement,<sup>9</sup> while others have found the effectiveness of CA and braces to be similar.<sup>10</sup> However, CA is advantageous in giving superior aesthetics, greater comfort and an easier oral hygiene.<sup>11</sup> The majority of the studies involving a comparative group, has significantly older patients in the group treated with Invisalign<sup>\*</sup>,<sup>12</sup> this has resulted in the alteration of movement or of groups of teeth.

# 2 Material and methods

### 2.1 Clinical examination and diagnosis

Ten consecutive healthy (ASA 1) patients (7 females and 3 males, between the ages of 17–19, average 18,3y) were treated. The intraoral examination revealed no periodontal problems, no active carious lesions and no TMJ problems. The extraoral examination showed a prognathic profile with lip incompetence at rest. All the patients had dental and skeletal Class III malocclusion with moderate crowding in both dental arches, negative overjet and the protrusion of the upper incisors and the retrusion of the lower incisors (dental compensation). The cephalometric analysis at T0 indicated a skeletal Class III and a hyperdivergent pattern of the vertical dimension, the proclination of the upper incisors and the lingualization of the lower incisors (Table 1). Three-dimensional radiographic examinations with CBCT were taken before treatment (T0) (Fig. 1) and a year after the orthognathic surgery (T1) (Fig. 2). From this analysis, the lingual or labial inclinations for each individual tooth/root were evaluated by using the axial slices perpendicular to its long axis. The image slices, perpendicular to the axial ones, were automatically reconstructed. This results in optimal visualization of the Marginal Bone Level (MBL) in relation to the Cemento-Enamel Junction (CEJ) in axial, coronal, and sagittal views, as described by Lund.<sup>1</sup> All the cases underwent a pre-orthognathic orthodontic treatment phase in order to correct dental compensation of the skeletal discrepancy. The patients were informed of the procedure and they or their parents, signed a consent form. Three-dimensional digital planning was performed with a dedicated software (ClinCheck<sup>®</sup>, Align Technology). The treatment was carried out with CA (Invisalign<sup>®</sup>, Align Technology). The RC (PAOO®) was performed 2-4 weeks into the orthodontic treatment, so as to regenerate bone in the direction of the dental movement, reducing the risks of long-term periodontal defects. The procedures and the materials used were conventional. The Code of Ethics of the World Medical Association (Declaration of Helsinki) was respected as was the approval by the ethical board of the Hospital.



(i) The table layout displayed in this section is not how it will appear in the final version. The representation below is solely purposed for providing corrections to the table. To preview the actual presentation of the table, please view the Proof.

Analysis of all the proclination of the lower incisors, in <u>10</u><sup>2</sup> cases, in each tooth, as decompensation of the Class III, programmed by the orthodontist and performed during the orthodontic treatment of preparation for maxillofacial surgery. In the first case, with <u>IMPA: 80,3° the planned proclination of lower incisors was +9</u>,7° (mean); in the second case with IMPA: 78,7° it was +9,975° (mean); in the third case with IMPA: 76,9° it was performed in +13° (mean).

Patient	Case1	Case2	Case3	Case4	Case5	Case6	Case7	Case8	Case9	Case10
IMPA T0 (°)	80,3	78,7	76,9	77	80.5	78.3	77.9	79.4	78.9	80.1
IMPA T1 (°)	87,5	87,7	88,1	84.3	89.6	88.2	87.3	89	88.7	89.2
Difference between the IMPA T1 and IMPA T0 (°)	(+7,2)	(+9)	(+11,2)	+7.3	+9.1	+9.9	+9.4	+9.6	+9.8	+9.1
Planned Proclination	+9.7	+9.2	12	8.3	+9.5	11	+10.8	+10.5	+10.7	+9.9
Difference between the planned and the obtained Proclination (°)	-2,5	-0,2	-0,8	-1	-0.4	-1.1	-1.4	-0.9	-0.9	-0.8
Difference between the programmed and the obtained of Proclination (%)	74.22	97.8	93.3	87.9	95.8	90	87	91.4	91.6	91.9

Legend: IMPA: angle between mandibular plane and inclination of lower incisors on lateral teleradiography.



Fig. 1



Pre-operative cephalometric showing a class III skeletal discrepancy.





Post-operative cephalometric showing surgical correction of the skeletal class III discrepancy.

#### 2.2 Three-dimensional digital planning

Intra-oral scanning was performed (iTero<sup>®</sup> Element 1, Align Technology). The Digital treatment planning included decompensation, in preparation for the surgery. The software enables individual tooth movement, thereby, giving a more precise control of the movements throughout the treatment<sup>13</sup> (Table 1).

#### 2.3 Regenerative Corticotomy

Shortly after starting the orthodontic treatment, augmentation corticotomy was performed. The surgery was carried out during aligner number 2 and 4 in the lower canine to canine area. A new tunnel approach facilitated the post-surgical recovery<sup>14</sup> by performing a few small vertical incisions. Following corticotomy an acellular dermal collagen matrix of animal origin (Mucoderm <sup>®</sup> -Botiss Biomaterials Zossen, Germany) was trimmed to allow sub-periosteal insertion. A xenograft was placed underneath the membrane once it was secured. The intra and the extra-oral photography and the CBCT were recorded at 1 year. No complications or adverse reactions were recorded.

## **3 Results**

#### 3.1 Radiographic evaluation

CBCT examinations were performed before starting the orthodontic treatment and at the end of the treatment. All the examinations were made using a 9000 3D CBCT (Carestream Health, USA) unit, equipped with a flat-panel detector. The exposed volume was 50 mm by 30 mm (voxel size =  $0.679 \ \mu$  to  $0.2 \ mm$ , depending if a "stitching" of 3 consecutive volumes was performed to represent the entire jaw), encompassing the teeth in the jaw where corticotomy was carried out. Exposure parameters were: 70 kV, 8–10 mA (based on the subject's size), and a single 360° 24–72 s exposure time comprising a range of 235–468 projections. CBCT were performed to evaluate the thickness of bone and the 3D positioning of the roots in the alveolar ridge before treatment. Primary data reconstructions were made using the acquisition software (CS3D Imaging, Carestream Health, USA), resulting in perpendicular slices in axial, coronal, and sagittal planes of the image volume. Subsequently, a second reconstruction was made to obtain contiguous 0.5 mm thick slices.

The thickness of the buccal plate was evaluated at 3-5-7 mm from the CEJ on CBCT at T0 and T1 (the difference statistically significant P < 0.05) (Fig. 3). The mean difference was at the 3 mm level  $1.42 \pm 0.5$  mm, at the 5 mm  $1.98 \pm 0.66$  mm and at the 7 mm  $2.70 \pm 0.87$  mm (Table 2). The distance between the CEJ and BM was also evaluated, and a decrease of an average of  $5.5 \pm 3.2$  mm at T0 to  $1.39 \pm 0.53$  mm at T1 was recorded (the difference statistically significant (<0.05) (Fig. 4). The changes were statistically significant with P < 0.05 (Table 3). The mean proclination based on IMPA values was +9.16 +-1.19°. *No signs of apical resorptions were noticed.* 

alt-text: Fig. 3

Fig. 3



Pre- and post-operative measurements on CBCT sections. Thickness of the buccal plate were measured at 3 different level (3, 5 and 7 mm from CEJ) and compared. See Table 2 for Results.

lt-text: Table 2 Table 2					
<i>i</i> The table layout dis purposed for provic	played in this section is not how i ling corrections to the table. To p	it will appear in the review the actual p	e final version. presentation o	The representat of the table, pleas	tion below is solely se view the Proof.
Previous version					<u>Exp</u>
Buccal plate thickness at 2	3, 5 and 7 mm from CEJ (Cemento	o-Enamel Junction).	).		
lorizontal 3 mm Number of teeth (n) Averag			ckness	Standard I	Deviation
~		A 4 A		^ <b>^</b> ^	
Buccal plate thickness at 1	3, 5 and 7 mm from CEJ (Cemento	o-Enamel Junction).	).		
Buccal plate thickness at 2	3, 5 and 7 mm from CEJ (Cemento	o-Enamel Junction) N to	). Number of reeth (n)	Average Thickness	Standard Deviation
Buccal plate thickness at a Horizontal 3-mm	3, 5 and 7 mm from CEJ (Cemento ge the colour. Should be same as th	o-Enamel Junction) N te	). Number of reeth (n)	Average Thickness	Standard Deviation
Buccal plate thickness at a Horizontal 3-mm [Instruction: Can't chang table] <u>Horizontal 3 mm</u>	3, 5 and 7 mm from CEJ (Cemento ge the colour. Should be same as th	ne rst ofthe	). Number of eeth (n)	Average Thickness	Standard Deviation
Buccal plate thickness at 2 Horizontal 3-mm [Instruction: Can't chang table] <u>Horizontal 3 mm</u> Pre op	3, 5 and 7 mm from CEJ (Cemento	ne rst ofthe 4	). Number of eeth (n) 40	Average Thickness 0.19	Standard Deviation
Buccal plate thickness at 2 Horizontal 3-mm [Instruction: Can't chang table]Horizontal 3 mm Pre op Post op	3, 5 and 7 mm from CEJ (Cemento	D-Enamel Junction) N ta ne rst ofthe 4	). Number of eeth (n) 40	Average Thickness 0.19 1.65	Standard Deviation
Buccal plate thickness at 2 Horizontal 3-mm [Instruction: Can't chang table]Horizontal 3 mm Pre op Post op Horizontal 5 mm	3, 5 and 7 mm from CEJ (Cemento	o-Enamel Junction). N ta ne rst ofthe 4	). Number of eeth (n) 40	Average Thickness 0.19 1.65	Standard Deviation 0.30 0.60
Buccal plate thickness at 2 Horizontal 3-mm [Instruction: Can't chang table]Horizontal 3 mm Pre op Post op Horizontal 5 mm Pre op	3, 5 and 7 mm from CEJ (Cemento	D-Enamel Junction) N ta ne rst ofthe 4 4	). <b>Number of</b> <b>eeth (n)</b> 40 40 40	Average Thickness     0.19     1.65     0.29	Standard     Deviation     0.30     0.60     0.42
Buccal plate thickness at 2 Horizontal 3-mm [Instruction: Can't chang table]Horizontal 3 mm Pre op Post op Horizontal 5 mm Pre op Post op	3, 5 and 7 mm from CEJ (Cemento	D-Enamel Junction) N ta he rst ofthe 4 4 4 4 4 4 4	). Number of eeth (n) 40 40 40	Average Thickness     0.19     1.65     0.29     2.42	Standard Deviation   Deviation 0.30   0.30 0.60   0.42 0.83
Buccal plate thickness at 2 Horizontal 3-mm [Instruction: Can't change table]Horizontal 3 mm Pre op Post op Post op Post op Horizontal 5 mm Pre op Post op Horizontal 7 mm	3, 5 and 7 mm from CEJ (Cemento	D-Enamel Junction) N ta ne rst ofthe 4 4 4 4 4 4	). Number of eeth (n) 40 40 40	Average Thickness     0.19     1.65     0.29     2.42	Standard   Deviation   0.30   0.60   0.42   0.83
Buccal plate thickness at 2 Horizontal 3-mm [Instruction: Can't change table]Horizontal 3 mm Pre op Post op Horizontal 5 mm Pre op Post op Horizontal 7 mm Pre op	3, 5 and 7 mm from CEJ (Cemento	D-Enamel Junction). N ta he rst ofthe 4 4 4 4 4 4 4 4 4 4 4 4 4	). Number of eeth (n) 40 40 40 40 40 40	Average   Thickness   0.19   1.65   0.29   2.42   0.20   0.40	Standard Deviation   Deviation 0.30   0.30 0.60   0.42 0.83   0.83 0.60

Difference are statistically significant p < 0.05.





Pre- and post-operative measurements on CBCT sections. Distance from Cemento-Enamel Junction (CEJ) to Marginal Bone Level (BML) were compared. See Table 2 for Results.

### alt-text: Table 3

#### Table 3

(*i*) The table layout displayed in this section is not how it will appear in the final version. The representation below is solely purposed for providing corrections to the table. To preview the actual presentation of the table, please view the Proof.

Bone vertical height. Distance from CEJ (Cemento-Enamel Junction) to marginal bone level (MBL).

Vertical (CEJ MBL)	Number of teeth (n)	Average distance	Standard Deviation			
Pre op	40	5.40	3.20			
Post op	40	1.39	1.14			
)ifference are statistically significant with $n < 0.05$						

Difference are statistically significant with p < 0.05.

### 3.2 Soft tissue evaluation

The changes of the keratinized gingiva were evaluated comparing the intraoral pictures. The measurements were taken with an open source image-processing program (J Image, <u>https://imagej.net/</u>). The width measurements of the central upper right incisor form the STL file of each patient was recorded and used to calibrate the values (Fig. 5). The measurements were calculated at T0 and T1 slides for each single tooth. The keratinized gingiva width averaged  $1.42 \pm 0.36$  mm at T0 and  $4.16 \pm 2.25$  mm at T1 (statistically significant with P < 0.05) (Table 4).





Pre-operative view of soft tissue in the anterior inferior area (incisors).



### **3.3 Proclination values**

The proclination of the incisors after decompensation was compared to the values that were digitally programmed by the software. The mean proclination of the incisors compared to the digitally planned was -  $1 + -0.6^{\circ}$  (89.87 + -6.46%).

# **4 Discussion**

Gingival recession in orthodontic treatment is a growing concern.<sup>15</sup> A higher incidence of gingival recession in patients treated for transverse discrepancy has been observed,<sup>16</sup> authors have failed to correlate the two.<sup>17</sup> One hypothesis is that orthodontics may create marginal bone resorption thereby, weakening the anatomical site which may require time

to develop a recession.<sup>18</sup> Recessions are twelve times more likely to occur when more than 5 mm of crowding are present,.<sup>19</sup> A recent study<sup>20</sup> states that six years after treatment, the recession of the lower incisors is 8 times more likely to occur. Time and an inflammatory process may be necessary for soft tissue migration and gingival recession. It was retained that any expansive movement may cause bony dehiscence and eventually gingival recession.<sup>21</sup> Many studies have used two-dimensional radiography, restricting the safe expansive movement to proximal bone surfaces.<sup>22</sup> Recently, 3D radiographic analysis shows that Orthodontic movement may inadvertently reposition the teeth beyond the bony alveolar housing. This may create bone dehiscences and fenestrations.<sup>23</sup> The correlation between rapid palatal expansion and the thinning of the buccal bone was also demonstrated.<sup>21</sup>

PAOO<sup>®</sup> may help to modify the alveolar anatomy following the position of the roots.<sup>25</sup> Orthodontic treatment should aim to three-dimensionally position the roots inside the bony envelope at the end of the treatment.<sup>26</sup> The combination of corticotomy and a regenerative procedure has the ability to augment the original anatomy despite unfavorable root movement.<sup>27, 28</sup> A very recent Best Evidence Review of the American Academy of Periodontology confirmed that bone grafting and corticotomy, together with perfectly planned orthodontic treatment, may provide clinical benefits such as modifying periodontal phenotype, maintaining or enhancing buccal bone thickness, accelerating tooth movement, expanding the scope of safe tooth movement for patients undergoing orthodontic tooth movement.<sup>29</sup> In a retrospective analysis, this only happened when a bone regenerative procedure was combined with corticotomy. Whenever corticotomy was performed alone, the preservation of the existing bone volume was rarely achieved.<sup>8</sup> Despite the tunnel approach, it was possible to control the positioning of the grafting material and to regenerate bone, even in the most coronal portion of the alveolar bone, which is typically the most critical part. The T1 images of the CBCT in fact show two important findings: 1- the bone graft was precisely positioned at the marginal bone level; 2- the buccal plate massively increased despite the proclination of approximatively 10° compared to the T0 scans, both in thickness and in vertical height. This may clinically translate into more bone structure that correctly accommodates all the teeth within the alveolar bony envelope and therefore, prevents extractions of premolars, which is often contemplated in orthognathic cases.

# **5** Conclusions

The combination of CA orthodontic technique and RC seems to be an effective method to increase both soft and hard alveolar tissues in decompensation. This may happen despite an unfavorable movement of the roots outside the original bone anatomy. The tunnel approach, if correctly used, does not interfere with precise positioning of the graft. The association of 3-D digital planning and CA allows for an accurate control of the orthodontic movements. The use of the digital dental movements' software generates a treatment plan with details, both in terms of degrees and millimeters, with high orthodontic predictability. The pre-surgical decompensation with bone regeneration in skeletal Class III may avoid non-reversible treatment such as premolar extractions while lowering the risks of detrimental side effects on the periodontal structures.

# **DE Uncited References**

### 24

# Acknowledgements

28 Augmentation Material was provided by Botiss Biomaterials Zossen, Germany.

# References

(i) The corrections made in this section will be reviewed and approved by a journal production editor. The newly added/removed references and its citations will be reordered and rearranged by the production team.

- Lund H., Grondahl K., Grondahl H.G. Cone beam computed tomography evaluations of marginal alveolar bone before and after orthodontic treatment combined with premolar extractions. Eur J Oral Sci. 2012;120:201.
- 2. Choi Y.J., Chung C.J., Kim K.H. Angle Orthod. 2015 May;85(3):427-433.

- Renkema A.M., Navratilova Z., Mazurova K., Katsaros C., Fudalej P.S. Gingival labial recessions and the post-treatment proclination of mandibular incisors. Eur J Orthod. 2015 Oct;37(5):508–513.
- Wilcko M.T., Wilcko W.M., Pulver J.J., Bissada N.F., Bouquot J.E. Accelerated osteogenic orthodontics technique: a 1-stage surgically facilitated rapid orthodontic technique with alveolar augmentation. J Oral Maxillofac Surg. 2009;67:2149–2159.
- Brugnami F., Caiazzo A. Orthodontically driven corticotomy: tissue engineering to enhance orthodontic and multi-disciplinary treatment. Wiley – Blackwell; 2014.
- Wang B., Shen G., Fang B., Yu H., Wu Y. Corticotomy-assisted presurgical orthodontics of class III malocclusions: a cephalometric and cone-beam computed tomography study. J Craniofac Surg. 2013;24(6):1886–1890.
- Ahn H.W., Lee D.Y., Park Y.G., Kim S.H., Chung K.R., Nelson G. Morphologic evaluation of dentoalveolar structures of mandibular anterior teeth during augmented corticotomy-assisted decompensation. Am J Orthod Dentofacial Orthop. 2016;150:659–669.
- Brugnami F., Caiazzo A., Mehra P. Can corticotomy (with or without bone grafting) expand the limits of safe orthodontic therapy? J Oral Biol Craniofac Res. 2017;8:1–6.
- 9. Phan X., Ling P.H. Clinical limitations of Invisalign. J Can Dent Assoc. 2007;73:263–266.
- Zheng M., Liu R., Ni Z., Yu Z. Efficiency, effectiveness and treatment stability of clear aligners: a systematic review and meta-analysis. Orthod Craniofac Res. 2017;20:127–133.
- 11. Boyd R.L. Esthetic orthodontic treatment using the Invisalign appliance for moderate to complex malocclusions. J Dent Educ. 2008;72:948–967.
- Kuo E., Miller R.J. Automated custom-manufacturing technology in orthodontics. Am J Orthod Dentofacial Orthop. 2003;123(5):578–581.
- Rossini G., Parrini S., Castroflorio T., Deregibus A., Debernardi C.L. Efficacy of clear aligners in controlling orthodontic tooth movement: a systematic review. Angle Orthod. 2015;85(5):881–889.
- 14. Brugnami F., Caiazzo A., Mehra P. Tunnel regenerative corticotomy TURC. A new surgical approach for surgically facilitated orthodontics. A technical note. J Oral Biol Craniofac Res. 2021;11(1):63–65.
- Wennstrom J.L., Lindhe J., Sinclair F., et al. Mucogingival therapy. Annual Periodontoogy. 1996;1:671– 701.
- Graber L.W., Vanarsdall R.L., Vig K.W.L. Orthodontics: Current Principles & Techniques 4th Edition, Mosb. 2005.
- 17. Bassarelli T., Dalstra M., Melsen B. Changes in clinical crown height as a result of transverse expansion of the maxilla in adults. EJO (Eur J Orthod). 2005;27(2):121–128.
- Wennström J.L., Lindhe J., Sinclair F., Thilander B. Some periodontal tissue reactions to orthodontic tooth movement in monkeys. J Clin Periodontol. 1987;14:121–129.
- 19. Staufer K., Landmeser H. Effects of crowding in the lower anterior segment—a risk evaluation depending upon the degree of crowding. J Orofac Orthop. 2004;65:13.
- Renkema A.M., Fudalej P.S., Renkema A.A., et al. Gingival labial recessions in orthodontically treated and untreated individuals: a case-control study. Am J Orthod Dentofacial Orthop. 2013;143(2):206–212.
- 21. Engelking G., Zachrisson B.U. Effects of incisor repositioning on monkey periodontium after expansion through the cortical plate. Am J Orthoped. 1982;82:23–32.
- Hollender L., Ronnerman A., Thilander B. Root resorption, marginal bone support and clinical crown length in orthodontically treated patients. EJO (Eur J Orthod). 1980;2:197–205.

- 23. Sarikaya S., Haydar B., Ci?er S., et al. Changes in alveolar bone thickness due to retraction of anterior teeth. Am J Orthod Dentofacial Orthop. 2002;122(1):15–26.
- 24. Garib D.G., Henriques J.F., Janson G., et al. Periodontal effects of rapid maxillary expansion with toothtissue borne and tooth-borne expanders: a computed tomography evaluation. Am J Orthod Dentofacial Orthop. 2006;126:749–758.
- 25. Williams M.O., Murphy N.C. Beyond the ligament: a whole-bone periodontal view of dentofacial orthopedics and falsification of universal alveolar immutability. Semin Orthod. 2008;14:246–259.
- Fuhrmann R. Three-dimensional interpretation of periodontal lesions and remodeling during orthodontic treatment. Part III. J Orofac Orthop. 1996;57:224–237.
- 27. Brugnami F., Caiazzo A., Mehra P. Can corticotomy (with or without bone grafting) expand the limits of safe orthodontic therapy? J Oral Biol Craniofac Res. 2017;8(1):1–6.
- 28. Brugnami F, Caiazzo A, Signorelli L., Capri D. Regenerative orthodonties: GBR and corticotomy to stretch the limits of orthodontic treatment. Int J Periodontics Restor Dent. Accepted for publication.Brugnami F, Caiazzo A, Signorelli L., Capri D. Regenerative orthodontics: GBR and corticotomy to stretch the limits of orthodontic treatment. Int J Periodontics Restorative Dent. 2021 Jan-Feb;41(1):105-111. doi: 10.11607/prd.4562. PMID: 33528458
  - 29. Wang C.W., Yu S.H., Mandelaris G.A., Wang H.L. Is periodontal phenotype modification therapy beneficial for patients receiving orthodontic treatment? An American Academy of Periodontology best evidence review. J Periodontol. 2020;91:299–310.

### **Queries and Answers**

### Q1

Q4

Query: Please confirm that the provided email "fbrugnami@gmail.com" is the correct address for official communication, else provide an alternate e-mail address to replace the existing one, because private e-mail addresses should not be used in articles as the address for communication.

Answer: yes

### Q2

Query: Please check that the affiliations link the authors with their correct departments, institutions, and locations, and correct if necessary.

Answer: Correspnding author address is "Piazza dei Prati degli Strozzi 21" and not "Piazza dei Prati degli Strozi 21"

#### Q3

Query: Correctly acknowledging the primary funders and grant IDs of your research is important to ensure compliance with funder policies. We could not find any acknowledgement of funding sources in your text. Is this correct? Answer: Yes Query: Please update Ref. 28.

Answer: Refrence updated

#### Q5

**Query:** Uncited references: This section comprises references that occur in the reference list but not in the body of the text. Please position each reference in the text or, alternatively, delete it. Any reference not dealt with will be retained in this section. Thank you.

Answer: 24 is after 23 and before 25. by mistake indicated as 21.

### Q6

**Query:** Please confirm that given names and surnames have been identified correctly and are presented in the desired order and please carefully verify the spelling of all authors' names.

Answer: Yes